FREE NOTES · NCERT + JEE LEVEL

Integrals

Integration yaani anti-derivative, standard formulas, substitution, trigonometric identities, partial fractions, integration by parts, special integrals, definite integrals, fundamental theorem aur definite integrals ki properties.

01 Integration yaani anti-derivative
02 Standard formulas
03 Substitution se integration
04 Partial fractions
05 Integration by parts
06 Special integrals
07 Definite integrals & properties
08 JEE Extra + Tricks
09 Solved examples
10 Common mistakes
11 Practice questions
12 Quick revision: exam se 5 minute pehle

01 Integration yaani anti-derivative

Agar d/dx F(x) = f(x), to ∫f(x) dx = F(x) + C (indefinite integral; C = constant of integration). ∫[f ± g] = ∫f ± ∫g; ∫kf = k∫f.

02 Standard formulas

$\int f(x)\,dx$	Result	$\int f(x)\,dx$	Result
x^n $(n \neq -1)$	$x^{n+1}/(n+1)$	$1/x$	$\log\|x\|$
e^x	e^x	a^x	$a^x/\log a$
$\sin x$	$-\cos x$	$\cos x$	$\sin x$
$\sec^2 x$	$\tan x$	$\operatorname{cosec}^2 x$	$-\cot x$
$\sec x \tan x$	$\sec x$	$\operatorname{cosec} x \cot x$	$-\operatorname{cosec} x$
$\tan x$	$\log\|\sec x\|$	$\cot x$	$\log\|\sin x\|$
$\sec x$	$\log\|\sec x + \tan x\|$	$\operatorname{cosec} x$	$\log\|\operatorname{cosec} x - \cot x\|$
$1/\sqrt{(1 - x^2)}$	$\sin^{-1} x$	$1/(1 + x^2)$	$\tan^{-1} x$

Har indefinite integral me + C lagao.

03 Substitution se integration

Andar wale function ko t rakho taaki uska derivative bhi maujood ho. $\int f(g(x))g'(x)dx = \int f(t)dt$.

- $\int 2x \sin(x^2)\,dx$: $t = x^2 \Rightarrow -\cos(x^2) + C$.
- $\int f'(x)/f(x)\,dx = \log|f(x)| + C$.
- Trig powers: identities lagao ($\sin^2 x = (1 - \cos 2x)/2$, $\sin^3 x = (3 \sin x - \sin 3x)/4$, guna → jod).

04 Partial fractions

Denominator ka factor	Partial fraction
$(x - a)(x - b)$	$A/(x - a) + B/(x - b)$
$(x - a)^2$	$A/(x - a) + B/(x - a)^2$
$(x - a)(x^2 + bx + c)$	$A/(x - a) + (Bx + C)/(x^2 + bx + c)$

Numerator ki degree ≥ denominator ho to pehle bhaag do.

05 Integration by parts

$$\int u\,v\,dx = u\int v\,dx - \int[u' \int v\,dx]\,dx$$

u **ILATE** se chuno: Inverse trig, Log, Algebraic, Trig, Exponential (list me pehla = u). $\int e^x[f(x) + f'(x)]\,dx = e^x f(x) + C$.

Example: $\int x e^x\,dx = x e^x - e^x + C$. $\int \log x\,dx = x \log x - x + C$.

06 Special integrals


$\int$	**Result**
$dx/(x^2 - a^2)$	$(1/2a) \log\lvert(x - a)/(x + a)\rvert$
$dx/(a^2 - x^2)$	$(1/2a) \log\lvert(a + x)/(a - x)\rvert$
$dx/(x^2 + a^2)$	$(1/a) \tan^{-1}(x/a)$
$dx/\sqrt{(x^2 - a^2)}$	$\log\lvert x + \sqrt{(x^2 - a^2)}\rvert$
$dx/\sqrt{(a^2 - x^2)}$	$\sin^{-1}(x/a)$
$dx/\sqrt{(x^2 + a^2)}$	$\log\lvert x + \sqrt{(x^2 + a^2)}\rvert$
$\sqrt{(a^2 - x^2)}\,dx$	$(x/2)\sqrt{(a^2 - x^2)} + (a^2/2) \sin^{-1}(x/a)$
$\sqrt{(x^2 \pm a^2)}\,dx$	$(x/2)\sqrt{(x^2 \pm a^2)} \pm (a^2/2) \log\lvert x + \sqrt{(x^2 \pm a^2)}\rvert$

$ax^2 + bx + c$ ke liye: square poora karo. $(px + q)/(ax^2 + bx + c)$ ke liye: $px + q = A \cdot d/dx(ax^2 + bx + c) + B$ likho.

07 Definite integrals & properties

$$\int_a^b f(x)\,dx = F(b) - F(a)$$

Second fundamental theorem. + C ki zaroorat nahi. a aur b ke beech curve ke neeche area (sign ke saath).



Definite integral sums ki limit ke roop me

Property	Statement
P1	$\int_a^b f = -\int_a^b f$; $\int_a^a f = 0$
P2	$\int_a^b f = \int_a^c f + \int_c^b f$
P3 (king)	$\int_a^b f(x)\,dx = \int_a^b f(a + b - x)\,dx$
P4	$\int_0^a f(x)\,dx = \int_0^a f(a - x)\,dx$
P5	$\int_{-a}^a f = 2\int_0^a f$ agar f even; 0 agar f odd
P6	$\int_0^{2a} f = 2\int_0^a f$ agar $f(2a - x) = f(x)$; 0 agar $f(2a - x) = -f(x)$

08 JEE Extra + Tricks

JEE LEVEL

- $\int e^x[f(x) + f'(x)]dx = e^x f(x) + C$ (bahut aam).
- $\int dx/(a + b\cos x)$ jaise: $t = \tan(x/2)$.
- King property: $\int_0^a f(x)dx = \int_0^a f(a - x)dx \Rightarrow \int_0^{\pi/2} \sin^n x/(\sin^n x + \cos^n x)\, dx = \pi/4$.
- $\int_{-a}^{a} f = 2\int_0^a f$ (even), 0 (odd); periodic: $\int_0^{nT} f = n\int_0^T f$.
- $\int_0^{\pi/2} \sin^n x\, dx$ ke liye Walli: $[(n - 1)(n - 3)\ldots]/[n(n - 2)\ldots] \times$ (n even ho to $\pi/2$).
- By parts me ILATE: Inverse, Log, Algebraic, Trig, Exponential (pehla function).
- Leibniz rule: $d/dx \int_{g(x)}^{h(x)} f(t)dt = f(h)h' - f(g)g'$.

09 Solved examples

Q1. $\int (x^3 + 5x^2 - 4)/x^2\, dx$.

$\int(x + 5 - 4x^{-2})\, dx = x^2/2 + 5x + 4/x + C$.

Q2. $\int dx/((x + 1)(x + 2))$.

$1/(x+1) - 1/(x+2) \Rightarrow \log|(x + 1)/(x + 2)| + C$.

Q3. $\int x \cos x\, dx$.

$u = x$: $x \sin x - \int \sin x\, dx = x \sin x + \cos x + C$.

Q4. $\int_0^{\pi/2} \sqrt{(\sin x)}/(\sqrt{(\sin x)} + \sqrt{(\cos x)})\, dx$.

P4 se $I = \int \sqrt{(\cos x)}/(\sqrt{\cos} + \sqrt{\sin}) \Rightarrow 2I = \int_0^{\pi/2} 1\, dx = \pi/2 \Rightarrow$ **$I = \pi/4$**.

Q5. $\int_{-1}^{1} x^5 \cos x\, dx$ aur $\int dx/(x^2 + 2x + 2)$ nikalo.

Odd function $\Rightarrow$ 0. Doosra: $(x + 1)^2 + 1 \Rightarrow \tan^{-1}(x + 1) + C$.

10 Common mistakes

- Indefinite integral me + C bhool jaana.
- Definite integral me substitution ke baad limits na badalna.
- By parts me u galat chunna (ILATE follow karo).
- ∫ 1/x dx = log x likhna (log|x| hona chahiye).

11 Practice questions

1. ∫ $\sin^2 x\,dx$.
2. ∫ $e^x(\sin x + \cos x)\,dx$.
3. $\int_0^1 x e^x\,dx$.
4. $\int_0^{\pi/2} \sin^2 x\,dx$.

ANSWERS

1) $x/2 - \sin 2x/4 + C$ · 2) $e^x \sin x + C$ · 3) 1 · 4) π/4

12 Quick revision: exam se 5 minute pehle

METHODS

- Substitution: andar wala = t.
- Rationals ke liye partial fractions.
- Parts: ILATE.

KEY RESULTS

- $f'/f \Rightarrow \log|f|$.
- $e^x(f + f') \Rightarrow e^x f$.

SPECIAL

- $1/(x^2+a^2) \Rightarrow (1/a)\tan^{-1}(x/a)$.
- Square poora karo.

DEFINITE

- F(b) − F(a).
- a + b − x trick; odd ⇒ 0.

Koi doubt? padhaiadda.netlify.app pe AI Doubt Solver se poocho, ya is chapter ka mock test do.